

Treatment of H. Pylori Infections in Peptic Ulcer Disease

Introduction to H. Pylori

- Helicobacter Pylori, or H. Pylori is a bacterium that may cause ulcers in the stomach.
- H. pylori is treated with a course of antibiotics. It is important to complete the course of medicines that are prescribed.
- Failure to complete your medications will result in persistent H. pylori infection in the stomach.



Your Medications

Your Antibiotics (Complete the course)

<input type="checkbox"/> Amoxicillin 250mg Capsule	<ul style="list-style-type: none">• Take ___ capsules, 2 times a day• To be taken with or after food• Common Side Effects: Stomach discomfort, nausea, vomiting and diarrhoea
<input type="checkbox"/> Clarithromycin 500mg Tablet	<ul style="list-style-type: none">• Take ___ tablets, 2 times a day• To be taken with or after food• Common Side Effects: Nausea, vomiting and diarrhoea
<input type="checkbox"/> Metronidazole 200mg Tablet	<ul style="list-style-type: none">• Take 2 tablets, 3 times a day• To be taken with or after food• Common Side Effects: Metallic taste, dizziness, headache, diarrhoea, constipation, nausea, vomiting and abdominal cramps• Note: Avoid alcohol intake while on this medication.
<input type="checkbox"/> Tetracycline 250mg Capsule	<ul style="list-style-type: none">• Take 2 capsules, ___ times a day• To be taken 1 hour before, or 2 hours after food with a full glass of water• Common Side Effects: Diarrhoea, nausea, vomiting and increased skin sensitivity to light• Note: Avoid prolonged exposure to sunlight while on this medication. Space apart antacids, dairy products, calcium and iron supplements, 2 hours apart from this medication.


Your Medications

- Omeprazole 20mg Capsule**
 - Take ___ capsules, 2 times a day
 - To be taken 30 minutes to 1 hour before food
 - Common side effects: Headache, upset stomach, diarrhoea and dizziness
- Esomeprazole 20mg Capsule**
 - Take ___ capsules, 2 times a day
 - To be taken 30 minutes to 1 hour before food
 - Common side effects: Headache, upset stomach, diarrhoea and dizziness
- Bismuth Subcitrate 120mg Tablet**
 - Take 1 tablet, 4 times a day
 - To be taken before food
 - Common side effects: Darkening of mouth, tongue, teeth, nausea, vomiting and grayish or black stools

Note: Space apart bismuth from antacids or milk 30 minutes to 1 hour after taking bismuth
- Vonoprazan 20mg Tablet**
 - Take 1 tablet, 2 times a day
 - May be taken with or without food
 - Common side effects: Diarrhoea, constipation, nausea, bloatedness and changes in taste

Should you develop symptoms such as itchiness, rashes on the skin, swelling around the eyes, or difficulty breathing, please stop these medications and immediately seek medical attention.

Please do not hesitate to contact Pharmacy should you have any further questions about your medications.



Contributed by Pharmacy, Allied Health Services

This brochure is produced for educational purposes and should not be used as a substitute for medical diagnosis or treatment. Please seek the advice of a qualified healthcare provider before starting any treatment or if you have any questions related to your health or medical condition.

Information shared is accurate as of April 2024 and subject to revision without prior notice.



Scan QR code to download e-brochure